

New Release!

16mm Cotton Tipped Swab with Chlorhexidine-Alcohol 1% Solution 1%CHG PUSH SWAB

1%CHG \cdot **80vol%ethanol solution** + **16mm cotton-tipped swab set**

1%CHG Push Swab consists of cotton tipped swab and 1% chlorhexidine-alcohol solution(contains 80% alcohol) packed in separate compartments respectively in the same pouch.

Laborsaving and **Efficient at disinfecting**

It is recommended by the infection control guidelines to use CHG-alcohol swab at the concentrations above 0.5%.


16mm swab × 1 stick/pouch dosage per pouch: 5.0mL




Contains

80vol% Alcoho






Hold the pouch upright and keep the stick into the liquid solution downside.



Push down the swab Open the seal at the inside the pouch.



upside.



Pull out swab stick from the pouch.

External Disinfectant

16mm Cotton Tipped Swab with Chlorhexidine-Alcohol 1% Solution 1%CHG PUSH SWAB

Cautions for Storage: keep in a light shielding airtight container and avoid contact with heat sources and fire sources. Expiry date is written on the package.

Prohibitions(Do not use on the following patients and parts)

- 1. Patients who are allergic to CHG or alcohol.
- 2. Brain, spinal cord, inner ear, middle ear and outer ear.
- 3. Mucous membrane of lumen, urinary bladder and oral cavity.
- 4. Open wounds and mucous membrane.
- 5. Eyes

Ingredients and Dosage

1.Ingredients

Main Ingredients	Each 100 mL contains 1g of chlorhexidine gluconate.
Additive	Ethanol, pH control chemical.
Base Material	Cotton tipped swab.
Dosage	5 mL of solution per packet.

2.Properties of solution

Colorless transparent liquid with characteristic odor.

Purpose of Use

To disinfect skin and fingers.

Usage

Apply the solution on skin or finger after rinsing the surface.

Precautions

- 1.Careful adiministration is required for the following patients.
- (1) Patients who are allergic to certain medication.
- (2) Patients with family history in asthma.
- 2.Important precautions
- (1) In order to predict reaction such as shock or anaphylaxis, a sufficient medical interview should be performed regarding the history of hypersensitivity to chlorhexidine preparations and for predisposition to drug hypersensitivity in advance.
- (2) Do not use this product to disinfect external genitalia and bladder.
- (3) Avoid contact with eyes. In case of contact, rinse eyes immediately.
- (4) Be aware of alcohol inhalation when applying this product for a long period of time or a wide surface of area. (Inhalation of alcohol vapor may cause headache and mucous membrane irritation.)

3. Adverse reactions

Treatment outcome study has not been conducted to specify the possibility of occurrence of adverse reactions.

- (1) Significant adverse reactions There is a risk of occurrence of anaphylactic shock when using this product. Be aware of occurrence of any negative effects like hypotension, hives, difficulty in breathing, etc. Stop using this product immediately and take proper action promptly if any of these symptoms occur.
- (2) Other adverse reactions
 Rash and irritated skin. Stop using this product immediately when any irritating reaction occurs.
- 4. Application notes
 - (1) For external use only.
 - (2) This product may be deactivated by soap. Make sure to rinse the soap residue off completely before appying this product.
 - (3) Skin may be degreased and become rough when applying this product on the same site repeatedly.
 - (4) Remove protein stains such as blood and pus before applying this product. protein substance may weaken disinfecting efficacy.
 - (5) Flamable. Keep away from fire.

Disinfecting Efficacy

Applicable organism: gram-positive bacteria, gram-negative bacteria, tubercle basillus, fungus, part of virus.

Handling Precautions

- 1.Do not bleach when the clothing is soiled by this product. It may cause brown stain.
- 2. Use immediately after opening the seal.

Manufacturer

HAKUZO MEDICAL CORPORATION 2-4-9, Tokuicho, Chuo-ku, Osaka, JAPAN 540-0025 TEL: +816-6809-4319 FAX: +816-6495-0539 MADE IN JAPAN



HAKUZO MEDICAL CORPORATION

2-4-9, Tokuicho, Chuo-ku, Osaka, JAPAN

F	Product No.	Spec.	Qty/box	Qty/case	Bar Code
	2600323	16mm cotton-tipped swab Solution: 5.0mL	24 sticks	16 boxes	(01)14958286074594



Add: 2-4-9, Tokuicho, Chuo-ku, Osaka 540-0025, Japan

TEL: 06-6942-0451 FAX: 06-6945-0539

www.hakuzo.co.jp/eng/index.php